

# WFS1 A180P — Wolframin

Alanine → Proline at position 180. N-terminal cytoplasmic (intrinsically disordered). ClinVar Uncertain significance, AlphaMissense 0.998, DynaMut2  $\Delta\Delta G$  +0.66 kcal/mol (stabilising).

## IDENTITY

Variant	A180P (p.Alanine180Proline)
DNA change	c.538G>C
Gene · Protein	WFS1 · Wolframin (890 aa)
UniProt	O76024 · WFS1_HUMAN
ClinVar accession	VCV004718918
Amino acid change	Alanine (A) → Proline (P)

## STRUCTURAL CONTEXT

AlphaFold model	AF-O76024-F1, v6
pLDDT at residue 180	<b>90.00</b> HIGH CONFIDENCE
Domain	N-terminal cytoplasmic (intrinsically disordered)
Position context	N-terminal cytoplasmic (intrinsically disordered)
IDR flag	No — pLDDT well above 50 threshold

Position 180 sits in N-terminal cytoplasmic (intrinsically disordered). The wild-type residue is small/hydrophobic (alanine — methyl sidechain); the mutant is rigid/helix-breaking (proline — kinks backbone). The chemistry shift implies altered local packing, hydrogen-bonding, and/or electrostatics at this site.

## COMPUTATIONAL PREDICTIONS

ALPHAMISSENSE

**0.998**am\_class: **likely pathogenic** —  
threshold > 0.564DYNAMUT2  $\Delta\Delta G$ **0.66** kcal/mol

Stabilising · Job 178090202722

PLDDT (ALPHAFOLD)

**90.00**

high confidence

## CLINICAL EVIDENCE

ClinVar classification	UNCERTAIN SIGNIFICANCE
Review status	criteria provided, single submitter
Last evaluated	2025/09/03 00:00
Inheritance	Inheritance pattern not specified in ClinVar entry; WFS1 has both AD and AR presentations.
WFS1 variant landscape	A180P is 1 of ~326 pathogenic-spectrum variants in WFS1 (out of 2,243 in ClinVar)
	<ul style="list-style-type: none"> <li>(no conditions catalogued)</li> </ul>

## RESEARCH PATH DECISION TREE

$\Delta\Delta G < 2$  + binding site affected → CATEGORY 3 – docking experiments  $\Delta\Delta G 2-4$  → CATEGORY 2 – pharmacological chaperones  $\Delta\Delta G > 4$  → CATEGORY 1 – gene therapy pLDDT < 50 → CATEGORY 5 – IDR, experimental only Stable fold + functional site hit → CATEGORY 4 – site-specific docking

### Category 3/4 — Most Druggable

$|\Delta\Delta G|=0.66 < 2$  kcal/mol (fold intact) + AlphaMissense 0.998 confirms functional impact. Specific local contacts disrupted — priority for docking and pharmacological chaperone screening.

Wolframin's fold survives this substitution ( $|\Delta\Delta G|=0.66$  kcal/mol). The pathogenic signal is real — AlphaMissense places it at 0.998. Protein still folds, but a specific local site is broken. Pharmacological chaperones and small-molecule binders are the rational therapeutic vector.