

WFS1 S551F — Wolframin

Serine → Phenylalanine at position 551. Transmembrane helix 8. ClinVar Uncertain significance, AlphaMissense 0.962, DynaMut2 $\Delta\Delta G$ -0.87 kcal/mol (destabilising).

IDENTITY

Variant	S551F (p.Serine551Phenylalanine)
DNA change	c.1652C>T
Gene · Protein	WFS1 · Wolframin (890 aa)
UniProt	O76024 · WFS1_HUMAN
ClinVar accession	VCV002110404
Amino acid change	Serine (S) → Phenylalanine (F)

STRUCTURAL CONTEXT

AlphaFold model	AF-O76024-F1, v6
pLDDT at residue 551	85.94 HIGH CONFIDENCE
Domain	Transmembrane helix 8
Position context	Inside Transmembrane helix 8 · position 551 is bilayer-embedded
IDR flag	No — pLDDT well above 50 threshold

Position 551 sits in a transmembrane helix (Transmembrane helix 8). Wolframin has eleven such helices anchoring it in the ER membrane; substitutions inside the bilayer-embedded segments can disrupt helix packing, lipid contacts, and the overall ER topology of the protein. The wild-type residue is small polar (serine — hydroxyl); the mutant is large aromatic hydrophobic (phenylalanine). The chemistry shift implies altered local packing, hydrogen-bonding, and/or electrostatics at this site.

COMPUTATIONAL PREDICTIONS

ALPHAMISSENSE

0.962am_class: **likely pathogenic** — threshold > 0.564DYNAMUT2 $\Delta\Delta G$ **-0.87** kcal/mol

Destabilising · Job 178092097867

PLDDT (ALPHAFOLD)

85.94

high confidence

CLINICAL EVIDENCE

ClinVar classification	UNCERTAIN SIGNIFICANCE
Review status	criteria provided, single submitter
Last evaluated	2025/02/03 00:00
Inheritance	Inheritance pattern not specified in ClinVar entry; WFS1 has both AD and AR presentations.
WFS1 variant landscape	S551F is 1 of ~326 pathogenic-spectrum variants in WFS1 (out of 2,243 in ClinVar) <ul style="list-style-type: none">(no conditions catalogued)

RESEARCH PATH DECISION TREE

$\Delta\Delta G < 2$ + binding site affected → CATEGORY 3 – docking experiments $\Delta\Delta G 2-4$ → CATEGORY 2 – pharmacological chaperones $\Delta\Delta G > 4$ → CATEGORY 1 – gene therapy pLDDT < 50 → CATEGORY 5 – IDR, experimental only Stable fold + functional site hit → CATEGORY 4 – site-specific docking

Category 3/4 — Most Druggable

$|\Delta\Delta G|=0.87 < 2$ kcal/mol (fold intact) + AlphaMissense 0.962 confirms functional impact. Specific local contacts disrupted — priority for docking and pharmacological chaperone screening.

Wolframín's fold survives this substitution ($|\Delta\Delta G|=0.87$ kcal/mol). The pathogenic signal is real — AlphaMissense places it at 0.962. Protein still folds, but a specific local site is broken. Pharmacological chaperones and small-molecule binders are the rational therapeutic vector.